Modelling the interactions between the hydrodynamics and the population dynamics of *Calanus finmarchicus* in the Gulf of St. Lawrence: (1) The "*Calanus* circuit".

Frédéric Maps (ISMER)


Pr. Bruno Zakardjian (LSEET-LEPI)

Dr. Stéphane Plourde (IML)

Pr. François Saucier (ISMER)











The Gulf of St. Lawrence (GSL) system :

- Semi-enclosed sea with contrasted topograpy and circulation features
- Seasonal sea-ice cover
- Estuarine-like general circulation, highly variable
- Productive pelagic ecosytem : C.finmarchicus









Currents, 1999

0 - 30m




Currents, 1999


30 - 100m



Currents, 1999

100 - Bottom





OBJECTIVES


(1) Test the few conceptual models regarding the persistence of the *C.finmarchicus* population in the GSL system

(Plourde & Runge 1993; Runge et al. 1999; Plourde et al. 2001; Zakardjian et al. 2003)

(2) Give a comprehensive picture of *C.finmarchicus* dynamics in the GSL system

MATHERIAL & METHODS








- Stage-based
- Transfer rate f (T)
- Egg production rate f (C), C = Diatoms + Flagellates (NPZD)
- Mortality rate f (N,T)
- Vertical migration


MATHERIAL & METHODS




MATHERIAL & METHODS

Biologic model Parametrization :


- Stage-based
- Transfer rate f (T)
- Egg production rate f (C), C = Diatoms + Flagellates (NPZD)
- Mortality rate f (N,T)
- Vertical migration
- Diapause window








Observed







Simulated - No migration








Simulated - Migration


























Simulated - Migration




April 1999 # females.m⁻²

August 1999 # females.m⁻²



Simulated - Migration




April 1999 # C4-5 & females.m⁻².s⁻¹

August 1999 # C4-5 & females.m⁻²

CONCLUSION

- Critical role of the swimming behaviour
- More mecanistic knowledge of the few key features of C.finmarchicus population dynamics is required:
 - Ontogenetic & nychtemeral migrations
 - Diapause Poster in Session 10, ID 3522
 - Mortality <--> Cannibalism Poster in Session 4, ID 3543



THANK YOU

- In situ data were provided by the Atlantic Zonal Monitoring Program (AZMP) of DFO Canada
- This work was partially funded by a NSERC grant to F.M.
- Attendance to this symposium was partly supported by the organisation of the 4th Zoooplakton Production Symposium
- Data analysis and visualization were performed with the Ferret program (NOAA – PMEL)