

REFERENCES

- Aasen, O. 1963. Length and growth of the porbeagle (*Lamna nasus*) in the north west Atlantic. Fisk. Dir. Skr. (Ser. Havunders) 13: 20-37.
- Angliss, R.P., DeMaster, D.P., and Lopez, A.L. 2001. Alaska marine mammal stock assessments, 2001. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-124. 203 p.
- Aydin, K.Y. 2000. Trophic feedback and carrying capacity of Pacific salmon (*Oncorhynchus* spp.) on the high seas of the Gulf of Alaska. Doctoral Dissertation, University of Washington, Seattle, 98195.
- Aydin, K.Y. and Friday, N. 2001. The early development of ECOSIM as a predictive multi-species fisheries management tool. Document SC/53/E3 presented to the IWC Scientific Committee July 2001 (unpublished). International Whaling Commission, Cambridge, UK. 8 p.
- Aydin, K.Y., Myers, K.W., and Walker, R.V. 2000. Variations in summer distribution of the prey of Pacific salmon (*Oncorhynchus* spp.) in the offshore Gulf of Alaska in relation to oceanographic conditions, 1994-98. N. Pac. Andr. Fish. Comm. Bull. 2: 43-54.
- Aydin, K.Y., Lapko, V.V., Radchenko, V.I., and Livingston, P.A. 2002. A comparison of the eastern and western Bering Sea shelf and slope ecosystems through the use of mass-balance food web models. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-130. 78 p.
- Balanov, A.A. and Radchenko, V.I. 1998. New data on the feeding and consumption behaviours of *Anotopterus pharao*. J. Ichthyol. 38(6): 447-453.
- Beamish, R.J. 1993. Climate and exceptional fish production off the west coast of North America. Can. J. Fish. Aquat. Sci. 50: 2270-2291.
- Beamish, R.J. 1999. An introduction to the PICES symposium on the ecosystem dynamics in the Eastern and Western Gyres of the subarctic Pacific. Prog. Oceanogr. 43: 157-161.
- Beamish, R.J., Kim, S., Terazaki, M., and Wooster, W.S. (Eds.) 1999a. Ecosystem Dynamics in the Eastern and Western Gyres of the Subarctic Pacific. Proceedings of the Science Board Symposium, North Pacific Marine Science Organization (PICES). Prog. Oceanogr. 43 (2-4): 157-487. Oxford: Pergamon Press, 1999.
- Beamish, R.J., Noakes, D.J., McFarlane, G.A., Klyashtorin, L., Ivanov, V.V., and Kurashov, V. 1999b. The regime concept and natural trends in the production of Pacific salmon. Can. J. Fish. Aquat. Sci. 56: 516-526.
- Beamish, R.J., Leask, K.D., Ivanov, O.A., Balanov, A.A., Orlov, A.M., and Sinclair, B. 1999c. The ecology, distribution, and abundance of midwater fishes of the Subarctic Pacific Gyres. Prog. Oceanogr. 43(2-4): 399-443.
- Beamish, R.J. and Mahnken, C. 2001. A critical size and period hypothesis to explain natural regulation of salmon abundance and the linkage to climate and climate change. Prog. Oceanogr. 49 (1-4): 423-437.
- Belyaev, V.A. and Ivanov, A.N. 1999. Dynamic processes in the fish community of the Northwest Pacific. M. Terazaki *et al.* (Eds.) Ecosystem dynamics of the Kuroshio-Oyashio Transition Region (Proc. Int. Mar. Sci. Symp., August 1998). 179-193.
- Bigg, M.A. 1990. Migration of northern fur seals (*Callorhinus ursinus*) off western North America. Can. Tech. Rep. Fish. Aquat. Sci. 1764. 64 p.
- Bigler, B.S., Welch, D.W., and Helle, J.H. 1996. A review of size trends among North Pacific salmon (*Oncorhynchus* spp.). Can. J. Fish. Aquat. Sci. 53: 455-465.
- Bonfil, R. 1994. Overview of the world elasmobranch fisheries. FAO Fish. Tech. Pap. FAO, Rome, Italy. 341: 119 p.
- Brodeur, R., McKinnell, S., Nagasawa, K., Percy, W., Radchenko, V., and Takagi, S. 1999. Epipelagic nekton of the North Pacific Subarctic and Transition Zones. Prog. Oceanogr. 43: 365-398.
- Brodeur, R.D. and Ware, D.M. 1995. Interdecadal variability in distribution and

- catch rates of epipelagic nekton in the Northeast Pacific Ocean. p. 329-356. In R.J. Beamish (Ed.) Climate change and northern fish populations, National Research Council of Canada, Ottawa.
- Buckland, S.T., Cattanach, K.L., and Miyashita, T. 1992. Minke whale abundance in the northwest Pacific and the Okhotsk Sea, estimated from 1989 and 1990 sighting surveys. Rep. Int. Whal. Commn. 42: 387-392.
- Buckland, S.T., Cattanach, K.L., and Hobbs, R.C. 1993. Abundance estimates of Pacific white-sided dolphin, northern right whale dolphin, Dall's porpoise and northern fur seal in the North Pacific, 1987-1990. Bulletin 53(III), North Pacific Commission. 387-407.
- Chapman, D.G. 1980. Estimates of pregnancy rates - North Pacific. Rep. Int. Whal. Commn. 2: 59-60.
- Chapman, D.G. 1973. Spawner-recruit models and estimation of the level of maximum sustainable catch. Rappports et proces-verbaux des reunions. 164: 325-332.
- Christensen, V., Walters, C.J., and Pauly, D. 2000. ECOPATH with ECOSIM: a Users Guide, October 2000 Edition. Fisheries Centre, University of British Columbia, Vancouver, Canada and ICLARM, Penang, Malaysia. 130 p.
- Christensen, V., and Pauly, D. 1992. ECOPATH II - a software for balancing steady-state ecosystem models and calculating network characteristics. Ecol. Modell. 61: 169-185.
- Clinton and LeBoeuf 1993. Sexual selection's effect on male life history and the pattern of male mortality. Ecology 74(6): 1884-1892
- Davis, N.D., Myers, K.W., and Ishida, Y. 1998. Caloric value of high-seas salmon prey organisms and simulated salmon ocean growth and prey consumption. N. Pac. Anadr. Fish. Comm. Bull. 1: 146-162.
- Doi, T., Nemoto, T., and Ohsumi, S. 1967. Memorandum on results of Japanese stock assessment of whales in the North Pacific. Rep. Int. Whal. Commn. 17: 111-115.
- Eslinger, D.L., Kashawai, M.B., Kishi, M.J., Megrey, B.A., Ware, D., and Werner, F.E. 2000. MODEL Task Team Workshop Report: Final Report of the International Workshop to Develop a Prototype Lower Trophic Level Ecosystem Model for Comparison of Different Marine Ecosystems in the North Pacific. PICES Scientific Report No. 15: 1-78.
- Evans, P. 1987. The natural history of whales and dolphins. Helm Publishers, London. 343 p.
- Faculty of Fisheries, Hokkaido University. 2000. Data Record of Oceanographic Observations and Exploratory Fishing No 43, 215pp.
- Falkowski, P.G., and Wilson, C. 1992. Phytoplankton productivity in the North Pacific ocean since 1900 and implications for absorption of anthropogenic CO₂. Nature. 358, 741-743.
- Favorite, F. and Laevastu, T. 1979. A study of the ocean migrations of sockeye salmon and estimation of the carrying-capacity of the North Pacific Ocean using a dynamical salmon ecosystem model (NOPASA). Northwest and Alaska Fisheries Center, NMFS, Seattle, WA. 47 p.
- Field, J. 2001 CalCOFI reports in press.
- Field, J.C, RC Francis and A Strom. 2001. Towards a fisheries ecosystem plan for the northern California current. CalCOFI Rep. 42: 74-87.
- Francis, R.C., Hare, S.R., Hollowed, A.B., and Wooster, W.S. 1998. Effects of interdecadal climate variability on the oceanic ecosystems of the NE Pacific. Fish. Oceanogr. 7: 1-21.
- Freeland, H., Denman, K., Wong, C.S., Whitney, F., and Jacques, R. 1998. Evidence of change in the winter mixed layer in the northeast Pacific Ocean. Deep-sea Res. I. 44: 2117-2129.
- Gambell, R. 1985a. Fin whale *Balaenoptera physalus* (Linneus, 1758). Handbook of Marine Mammals, Vol. 3. The sirenians and baleen whales: 171-192.
- Gambell, R. 1985b. Sei whale *Balaenoptera borealis* (Lesson, 1828). Handbook of Marine Mammals, Vol. 3. The sirenians and baleen whales: 155-170.
- Givens, G.H., Raftery, A.E., and Zeh, J.E. 1993. Benefits of a Bayesian approach for synthesizing multiple sources of evidence and uncertainty linked by a deterministic model. Rep. Int. Whal. Comm. 43: 495-500.
- Givens, G.H. 1994. A Bayesian framework and importance sampling methods for synthesizing multiple sources of evidence and uncertainty linked by a complex mechanistic model.

- Doctoral Dissertation, University of Washinton, Seattle, WA 98195. XII, 157p.
- Harrison, P.J., Boyd, P.W., Varela, D.E., Takeda, S., Shiimoto, A., and Odate, T. 1999. Comparison of factors controlling phytoplankton productivity in the NE and NW Subarctic Pacific Gyres. *Prog. Oceanog.* 43: 205-234.
- Hobbs, R.C. and Jones, L.L. 1993. Impacts of high seas driftnet fisheries on marine mammal populations in the North Pacific. *Bull. Int. N. Pac. Fish. Comm.* 53: 409-434.
- Hollowed, A.B., Hare, S.R., and Wooster, W.S. 1998. Pacific - basin climate variability and patterns of northeast Pacific marine fish production, pp. 89-104. *In* G. Holloway, P. Muller, and D. Henderson, (Eds.) Proceedings of the 10th 'Aha Huliko'a Hawaiian winter workshop on biotic impacts of extratropical climate variability in the Pacific, January 26-29, 1998. SOEST Special Publication.
- Horwood, J. 1990. Biology and exploitation of the Minke whale. CRC Press Inc. Boca Raton, FL.
- Hughes, S.E. 1964. Stock composition, growth, mortality and availability of Pacific saury *Cololabis saira*, of the northeastern Pacific ocean. *Fish. Bull.* 72(1): 121-131.
- Hunt, G.L., Kato, H., and McKinnell, S.M. (Eds.). 2000. Predation by marine birds and mammals in the subarctic North Pacific Ocean. PICES Scientific Report No. 14. 165 p.
- Ishida, Y., Ito S., Ueno Y., and Sakai, J. 1998. Seasonal growth patterns of Pacific salmon (*Oncorhynchus* spp.) in offshore waters of the North Pacific Ocean. *N. Pac. Anadr. Fish. Comm. Bull.* 1: 66-80.
- Ishida, Y., Ito, S., Kaeriyama, M., McKinnell, S., and Nagasawa, K. 1993. Recent changes in age and size of chum salmon (*Oncorhynchus keta*) in the North Pacific Ocean and possible causes. *Can. J. Fish. Aquat. Sci.* 50: 290-295.
- Ishida, Y., Ito, S., Kaeriyama, M., McKinnell, S., and Nagasawa, K. 1993 recent changes in age and size of chum salmon (*Oncorhynchus keta*) in the North Pacific Ocean and possible causes. *Can. J. Fish. Aquat. Sci.* 50: 290-295
- Ivanov O.A. 1997. Composition and biomass of fishes and cephalopods in upper mesopelagial layer in the Pacific waters off the Kuril Islands and Kamchatka. *Voprosy Ikhtiologii*, 37(2): 167-178 (in Russian)
- Kaeriyama, M., Nakamura, M., Yamaguchi, M., Ueda, H., Anma, G., Takagi, S., Aydin, K.Y., Walker, R.V., and Myers, K.W. 2000. Feeding ecology of sockeye and pink salmon in the Gulf of Alaska. *N. Pac. Anadr. Fish. Comm. Bull.* 2: 55-63.
- Kawamiya, M., Kishi, M., Yamanaka, Y., and Suginoara, N. 1997. Procuring reasonable results in different oceanic regimes with the same ecological-physical coupled model. *J. Oceanogr.* 53: 397-402.
- Kato, H., Miyashita, T., and Baba, N. 1997. Summary of population abundance estimates in the North Pacific based on Japanese data base. Documents submitted to WG11 meeting of PICES in Pusan. 12 p.
- Kuzin, A.E., Zolotukhin, S.F., Krupyanko, N.I., and Semenchenko, A. Yu. 2000. Incidental catch of the northern fur seal *Callorhinus ursinus* and the Dall's Porpoise *Phocoenoides dalli* with salmon gillnets in the Sea of Japan. *Russ. J. Mar. Biol.* 26 (2): 134-136.
- Longhurst, A., Sathyendranath, S., Platt, T., and Caverhill, C. 1995. An Estimate of global primary production in the ocean from satellite radiometer data. *Journal of Plankton Research.* 17: 1245-1271.
- Lapashina, V.I. 1988. On the feeding of the squid *Berryteuthis anonychus* Pearey and Voss from the area of the north-eastern Pacific. pp. 1-5, Deposited MS, Dep VINITI 25.10.88.
- Mackas, D.L., and Tsuda, A. 1999. Mesozooplankton in the eastern and western subarctic Pacific: community structure, seasonal life histories, and interannual variability. *Prog. Oceanog.* 43: 335-363.
- Mackay, A. 1981. The generalized inverse. *Practical Computing* (September): 108-110.
- Mantua, N.J., Hare, S.R., Zhang, Y., Wallace, J.M., and Francis, R.C. 1997. A Pacific interdecadal climate oscillation with impacts on salmon production. *Bull. Am. Meteorol. Soc.* 78: 1069-1080.
- McFarlane, G.A. and King, J.R. In Press. Migration patterns of spiny dogfish. *Fish. Bull.*
- McFarlane, G.A., Beamish, R.J., Yatsu A., and Krovnin A.S. 2001 BASS Workshop on the development of a conceptual model of the

- Subarctic Pacific basin ecosystems. PICES Sci. Rep. No.17, 1-24.
- Megrey, B.A., Kishi, M.J., Kashawai, M.B., Ware, D., Werner, F.E., Eslinger, D.L. .2000. PICES Lower Trophic Level Modeling Workshop, Nemuro 2000. *PICES Press* 8(2): 18-22.
- Melnikov, I. V. 1997. Pelagic fishes - consumers of Pacific salmon: distribution in the Russian economic zone and adjacent waters, abundance, and some biology features. *Izvestiya TINRO* 122: 213-238.
- Murata, M. and Shimazu, Y. 1982. On some population parameters of flying squid, *Ommastrephes bartrami* (Lesurur) in the North Pacific. *Bull. Hokkaido Reg. Fish. Res. Lab.* 47: 1-10.
- Myers, K.W., Aydin, K.Y., Walker, R.V., Fowler, S., and Dahlberg, M.L. 1996. Known ocean ranges of stocks of Pacific salmon and steelhead as shown by tagging experiments, 1956-1995. (NPAFC Doc. 192.). Univ. Washington, Fish. Res. Inst., Seattle.
- Ogi, H., Yatsu, A., Hatanaka, H., Nitta, A. 1993. The mortality of seabirds by driftnet fisheries in the North Pacific. *Bull. North. Pac. Commn.* 53 (3): 499-518.
- Ohsumi, S. 1979. Interspecies relationships among some biological parameters in cetaceans and estimation of the natural mortality coefficient of the southern hemisphere Minke whale. *Rep. Int. Whal. Commn.* 29: 397-406.
- Olesiuk, P.F., Bigg, M.A., and Ellis, G.M. 1990. Life history and population dynamics of resident killer whales (*Orcinus orca*) in the coastal waters of British Columbia and Washington State. *Rep. Int. Whal. Commn.* 2: 209-243
- Pauly, D. and Christensen, V. (Eds.). 1996. Mass-balance models of north-eastern Pacific ecosystems. Fisheries Centre, University of British Columbia. Research Reports 4. 131 p.
- Pauly, D., Christensen, V., and Walters, C.J. 2000. ECOPATH, ECOSIM and Ecospace as tools for evaluating ecosystem impact of fisheries. *ICES J. Mar. Sci.* 57: 697-706.
- Pauly, D., Trites, A.W., Capuli, E., and Christensen, V. 1998. Diet composition and trophic levels of marine mammals. *ICES J. Mar. Sci.* 55: 467-481.
- Pearcy, W.G., Fisher, J.P., Anma, G., and Meguro, T. 1996. Species associations of epipelagic nekton of the North Pacific Ocean, 1978-1993. *Fish. Oceanogr* 5: 1-20.
- Pearcy, W.G., Aydin, K.Y., and Brodeur, R.D. 1999. What is the carrying capacity of the North Pacific Ocean for salmonids? *PICES Press* 7(2): 17-23
- Pearcy, W.G., Brodeur, R.D., Shenker, J.M., Smoker, W.W., and Endo, Y. 1988. Food habits of Pacific salmon and steelhead trout, mid-water trawl catches and oceanographic conditions in the Gulf of Alaska, 1980-85. *Bull. Ocean. Res. Inst.* 26: 29-78.
- Polovina, J.J. 1985. An approach to estimating an ecosystem box model. *Fish Bull. US.* 83: 457-460.
- Polovina, J.J., Mitchum, G.T., and Evans, G.T. 1995. Decadal and basin-scale variation in mixed layer depth and the impact on biological production in the Central and North Pacific, 1960-88. *Deep-Sea Res.* 42: 1701-1716.
- Purcell, J. 1996. Lower trophic levels. UBC Fisheries Center Research Reports 4(1): 16.
- Ricker, W.E. 1995 Trends in the average size of Pacific salmon in Canadian catches. Pp 593-602. In: Beamish, R.J. [ed.] Climate change and northern fish populations. *Can. Spec. Publ. Fish. Aquat. Sci.* 121: 739 p.
- Rogers, D.E. 1999. Estimates of annual salmon runs from the North Pacific, 1951-1998. Fisheries Research Institute, Seattle, WA. 11 p.
- Sanger, G.A. 1972. Fishery potentials and estimated biological productivity of the subarctic Pacific region. pp. 561-574. In: *Biological Oceanography of the northern North Pacific Ocean*. A.Y. Takenouti (ed.). Idemitsu Shoten, Tokyo, 626 p.
- Savinykh, V.F. and Chuchukalo, V.I. 1996. Daily ration of the Pacific Pomfret, *Brama japonica*, in the Northwestern Pacific Ocean during feeding periods. *Russian J. Mar. Biol.* 22(6): 327-332.
- Shimizaki, K. 1986. Distribution of the pelagic fish community around the subarctic boundary in the north Pacific ocean. *INPFC* 47: 247-264.
- Shuntov V. P., Radchenko, V. I., Chuchukalo, V. I., Efimkin, A. Ya., Kuznetsova, N. A., Lapko,

- V. V., Poltev, Yu. N., and Senchenko, I. A. 1993a. Structure of planktonic and nektonic communities in the upper epipelagic zone in the western Bering Sea and Pacific Waters off Kamchatka in the period of anadromous migrations of salmon. *Russ. J. Mar. Biol.* 19(4): 231-239
- Shuntov V. P., Radchenko, V. I., Chuchukalo, V. I., Efimkin, A. Ya., Kuznetsova, N. A., Lapko, V. V., Poltev, Yu. N., and Senchenko, I. A. 1993b. Structure of planktonic and nektonic communities in the upper epipelagic zone of the Sakhalin-Kuril region in the period of anadromous migrations of salmon. *Russian J. Mar. Biol.* 19(4): 240-247
- Shuntov V. P., Lapko, V. V., Nadtpchiy, V. V., Samko, Ye. V. 1995. Annual changes in Ichthyocenes of the upper epipelagic zone of the Sakhalin-Kuril region. *Journal of Ichthyology.* 35(1): 32-43.
- Small, K.J. and DeMaster, D.P. 1995. Survival of five species of captive marine mammals. *Mar. Mamm. Sci.* 11(2): 209-226.
- Stewart, B., Yochem, P.K., Huber, H.R., DeLong, R.L., Jameson, R.J., Sydeman, W.J., Allen, S.G., and LeBoeuf, B.J. 1994. History and present status of the northern elephant seal population. pp. 29-48 In: Leboeuf, B. and R.H. Laws (Eds.). *Elephant seals: population ecology, behavior and psychology.* Univ. of California Press. 414 pp.
- Springer, A.M., Piatt, J.F., Shuntov, V.P., Van Vliet, G.B., Vladimirov, V.L., Kuzin, A.E., and Perlov, A.S. 1999. Marine birds and mammals of the Pacific Subarctic Gyres. *Prog. Oceanog.* 43: 443-487.
- Tadokoro, K., Ishida, Y., Davis, N.D., Ueyanagi, S., and Sugimoto, T. 1996. Change in chum salmon (*Oncorhynchus keta*) stomach contents associated with fluctuation of pink salmon (*O. gorbuscha*) abundance in the central subarctic Pacific and Bering Sea. *Fish. Oceanogr.* 5: 89-99.
- Tanaka, E. 1993. Population trajectories and effects of incidental take by driftnet on some dolphin populations using the Hitter/Fitter model. *INPFC Bull.* 53(1-3): 451-460.
- Taniguchi, A. 1999. Differences in the structure of the lower trophic levels of pelagic ecosystems in the eastern and western subarctic Pacific. *Prog. Oceanog.* 43: 289-315.
- Trites A. W., Livingston, P.A., Vasconcellos, M.C., Mackinson, S., Springer, A.M., and Pauly, D. 1999. Ecosystem change and the decline of marine mammals in the Eastern Bering Sea: testing the ecosystem shift and commercial whaling hypotheses. *Univ. British Columbia Fish. Cent. Rep.* 7(1): 106 p.
- TINRO. 1998a. Report on results of expedition aboard R/V "TINRO" in the Sea of Okhotsk in August-December 1998, Vladivostok, TINRO-Center, Vol. 2, 200 p. (in Russian).
- TINRO. 1998b. Report on results of expedition aboard R/V "TINRO" in the Sea of Okhotsk in August-December 1998, Vladivostok, TINRO-Center, Vol. 1, 446 p. (in Russian)
- Ulanowicz, R.E. 1983. Identifying the structure of cycling in ecosystems. *Math. Biosci.* 65: 219-237.
- Walters, C., Christensen, V., and Pauly, D. 1997. Structuring dynamic models of exploited ecosystems from trophic mass-balance assessments. *Rev. Fish. Biol. Fish.* 7: 139-172.
- Watanabe, Y., Oozeki, Y., and Kitagawa, D. 1997. Larval parameters determining preschooling juvenile production of Pacific saury (*Cololabis saira*) in the northwestern Pacific. *Can. J. Fish. Aquatic. Sci.* 54: 1067 - 1076.
- Wong, C.S., Whitney, F.A., Iseki, K., Page, J.S., and Zeng, J. 1995. Analysis of trends in primary production and chlorophyll-*a* over two decades at Ocean Station P (50°N, 145°W) in the subarctic Northeast Pacific Ocean, pp. 107-117, In: [ed.] R.J. Beamish, *Climate change and northern fish populations,* Can. Spec. Publ. Fish. Aquat. Sci. 121.