



Zooplankton response to NW Mediterranean hydroclimatic changes from 1966 to 2010

Pieter Vandromme¹, Lars **Stemmann**¹, Carmen García-Comas², Laure Mousseau¹, Franck Prejger¹, Ornella Passafiume¹, Marc Picheral³ and Gabriel Gorsky³

¹ Université Pierre et Marie Curie-Paris 6, UMR9073, Observatoire Océanologique de Villefranche sur Mer, BP 28, 06234 Villefranche sur Mer, France. E-mail: vandromme@obs-vlfr.fr

² Institute of Oceanography, National Taiwan University, No. 1, Sec. 4, Roosevelt Rd., Taipei 10617, Taiwan

³ CNRS UMR9073, Observatoire Océanologique de Villefranche sur Mer, BP 28, 06234 Villefranche sur Mer, France


The NW Mediterranean sea: CONTEXTE AND CHALLENGE

- functions like a mini ocean
 - change in anthropic input (notably N/P ratio) notably from Rhône river
 - changes in climate that change heat flux and water budget (1980's dry and 2000' dry), winter mixing and winter convection (**MEDOC** and **LIGURE**)


Bloom in Rhône river plume

Bloom in convection areas

The Ligurian bassin and the Point B sampling site are isolated from strong continental inputs.



1967-1993 Long term zooplankton time series at Point B



Shift and reorganization of zooplankton community at the end of the '80s (1987 shift year) toward an oligotrophic system.

What about the whole jellyfish, chaetognath and copepod populations?

Is oligotrophy confirmed after 17 more years of observations?

4 net collections since 1966



TS1: 1966-2011, Regent Net (680 μ m mesh), oblique hauls, daily sampling

TS2: 1966-2003, Juday Bogorov net (330 μ m mesh), vertical hauls, daily sampling

TS3: 1995–2006, WP2 net (200 μ m mesh), vertical hauls, weekly sampling

TS4: 2003-2011, WP2 net (200 μ m mesh) , vertical hauls, daily sampling


More than **20000** net samples but pooled in weekly jars (>**5000** jars).



Point B time series is one of longest and most complete
 → too many samples to analyse in a traditional way.

TS 2, 3 and 4 have been completely digitalized with the ZOOSCAN.

Automatic recognition followed by manual validation for all groups but copepods (TS1, TS2 and TS3)




Why collecting using 3 nets ?

TS1: REGENT

TS2: Juday
Bogorov

TS3 and 4, WP2



Different nets do not collect the same community

Most abundant organisms are small in the Med. sea.

BUT


plankton abundances not directly comparable.

Image analysis may provide a solution by measuring all organisms.


For more on combining nets results → March 16, W5, Automated visual plankton identification.

A warming Ligurian Sea


Surface Temperature (integrated 0-10m depth) from Point B




Earlier warming




Average year by decade




Warmer winters



Temperature at week 20



TS2: 1974-2003 zooplankton in JB nets



1st PC Zooplankton dataset (40%):

Copepod, Decapod, siphonophore, hydromedusae, chaetognathe


1974-1980: ↓ copepods and Jellies

1980-1990: ↑ All groups (phyto. 2 peaks)

1990-1998 ↓ All groups

1998-2003 ↑ All groups (phyto. 2 peaks)

(Carmen et al., in revision)



1st PC Environmental dataset (33%):

20,50m Temp. and Sal., atm.P, Ek.D., Rain, Sun


1974-1980: ↓ Salinity and Temp.

1980-1990: ↑ Salinity and Mean Temp .autumn

1990-1998 ↓ Salinity and ↑ Temp

1998-2003 ↑ Salinity


TS3: 1995-2005 zooplankton in WP2 nets



(Vandromme et al., in revision)



Cumulative sum of annual anomalies of the ten identified taxonomic groups.

Annual anomalies of total plankton biovolume and average seasonality of biovolume for the two periods.




TS1 : 1995-2005 zooplankton in Regent net

Cumulative sum of annual anomalies of two size classes of zooplankton in **Regent net** (Stemmann et al., in prep)




TS4 : 2004-2011 zooplankton in WP2

Only Copepoda



All but copepoda









- copepoda acartia
- copepoda calanoida
- copepoda candacia
- copepoda centropages
- copepoda harpact
- copepoda multiple
- copepoda oithona
- copepoda other
- copepoda poecilo corycaeus
- copepoda poecilo oncaea
- copepoda temora

- amphipods
- annelides
- appendicularia
- chaetognatha
- cladocera evadne
- cladocera penillia
- crust decapod
- crust decapoda large
- crust nauplii
- echinoderm
- egg
- fish
- gelatinous medusae
- gelatinous medusae ephyrales
- gelatinous thaliacae
- moll bivalve
- moll gasteropoda
- moll heteropoda
- moll pteropoda
- moll pteropoda cavolonia
- moll pteropoda creseis
- ostracods
- other
- radiolaria
- siphonophora


Zooplankton abundances have not declined since 2004

- the shift toward less zooplankton in late 80's observed by previous authors is confirmed.
- we observe a more recent shift in late 90's (confirmed in three independent nets) toward higher zooplankton concentration, therefore the “oligotrophisation” due to more stratification is not yet confirmed.
- in a warming Mediterranean Sea, what caused the different shifts ?
- can we extrapolate to the whole NW Mediterranean Sea ?


1995-2006: link with environmental data in WINTER (weeks 4-17)



Trend to more winter mixing and consequences for the ecosystem




Denser water in winter colder and saltier (close to intermediate Mediterranean water)







Less stratification in winter
→ more nutrient for
phytoplankton production and
zooplankton

More stratification in summer
→ better for jellyfish which feed
on more preys

Extrapolation to the whole NW Mediterranean sea



DYFAMED and Point B time series



(SYMPHONIE 3D model forced by ERA40 ciimatic model, Augier 2011)

- Zooplankton changes are closely linked to hydroclimate, and a bottom-up control is suggested (Comas et al., in revision) and in addition a top-down control of Zoo on Phyto (Vandromme et al., in revision).
- The stronger winter mixing in dry years (winter low T and high S) may be the principal factor to initiate the bloom and zooplankton development in spring despite the increase in summer stratification (and annual temperature).
- This may be a general pattern across the NW Mediterranean Sea.
- How long will that be ?